



The frequency of depressive mood and associated factors in pregnant women in a semi-urban region

Yarı kentsel bir bölgede gebe kadınlarda depresif duygudurum sıklığı ve ilişkili faktörler

Antepartum depressive mood and associated factors

Elif Yılmaz¹, Meltem Col², Yasemin Yavuz³

¹Department of Obstetrics and Gynecology, Dr.Sami Ulus Women's and Children's Health Teaching and Research Hospital, Sıhhiye,

²Department of Public Health, Ankara University Faculty of Medicine, Cebeci,

³Department of Biostatistics, Ankara University Faculty of Medicine, Cebeci, Ankara, Turkey

Öz

Amaç: Ülkemizin başkentinde yarı-kentsel bir bölgede gebelikte depresif duygu durum sıklığı ve ilgili risk faktörlerinin belirlenmesi. **Gereç ve Yöntem:** Çalışmaya gebeliğin son trimesterinde olan 600 gebe dahil edildi. Depresyon skorları "Edinburgh Depresyon Skalası" ile belirlendi. 13 ve üzerinde alanlar depresif kabul edildi. Depresif olan ve olmayan gebeler, sosyo-demografik özellikleri, ailesel ve medical özellikleri açısından karşılaştırıldı. **Bulgular:** Çalışmaya alınan altıyüz gebeden 153'ü depresif bulundu (25.5%). Düşük/ileri yaş, eşin/gebenin düşük eğitim seviyesi ve/veya işsiz olması, istem dışı yapılan evlilik, eşin akrabaları ile birlikte yaşama, planlanmamış gebelik ve kronik hastalık varlığı depresif duygu durum sıklığını arttıran faktörler olarak bulundu ($p<0.001$). **Tartışma:** Gebeliğin son trimesterinde depresif duygu durumun sık görüldüğü göz önünde tutulmalı ve depresyon taraması rutin gebelik takibi programlarının bir parçası haline getirilmelidir. Depresif duygu durum tanısı alan gebeler için de ileri tanı ve tedavi hizmetleri mutlaka sağlanmalıdır.

Anahtar Kelimeler

Gebelik; Depresif Duygu Durum; Risk Faktörleri

Abstract

Aim: To determine the antenatal depressive mood frequency and associated risk factors in a semi-urban region in the capital of our country. **Material and Method:** Six hundred pregnant women within their third trimester were included. Depression levels were evaluated according to the "Edinburgh Postnatal Depression Scale". Those scoring 13 and over were considered as depressed. Depressed and non-depressed pregnant women were compared according to socio-demographic characteristics, family, and medical features. **Results:** One hundred and fifty-three patients out of 600 were found to be depressive (25.5%). Low/advanced maternal age, low education level and unemployment of the women/husband, involuntary marriage, living with spouse's relatives, unplanned pregnancy, and existence of a diagnosed chronic disease were found to increase the frequency of depression ($p<0.001$). **Discussion:** Because of its high frequency among women in the third trimester, depression screening should be included in prenatal care routine programs and accessible diagnosis and treatment services should be provided for pregnant women diagnosed with depressive mood.

Keywords

Pregnancy; Depressive Mood; Risk Factors

DOI: 10.4328/JCAM.4972

Received: 12.03.2017 Accepted: 25.04.2017 Printed: 01.12.2017 J Clin Anal Med 2017;8(suppl 4): 261-5

Corresponding Author: Elif Yılmaz, Department of Obstetrics and Gynecology, Dr. Sami Ulus Women's and Children's Health Teaching and Research Hospital, 06200, Sıhhiye, Ankara, Turkey. T.: +90 3123056385 F.: +90 3123170353 E-Mail: elifkasyilmaz@gmail.com

Introduction

Depression is the most frequently observed psychological disorder during pregnancy and is an important public health issue because it is the most important risk factor for postnatal depression, it may also cause complications and negative fetal results during pregnancy, and its prevalence is even higher than in the postnatal phase [1,2]. Antenatal depression causes a reduction in mother's self-care and therefore may cause insufficient nutrition, alcohol-drug use, insufficient prenatal care, and even suicide attempts. Studies show that mothers experiencing depression during pregnancy have higher premature birth rates and give birth to infants with lower birth weights and lower APGAR scores [3,4]. It is also known that such mothers' children have a higher probability of developing behavioral and emotional problems/disorders in the future [5].

It has been reported that approximately 50% of women undergoing depression during pregnancy also suffered from postnatal depression [6]. There are many factors contributing to development of antenatal depression such as previously experienced depression attacks, marital problems, lack of spouse and social support, negative life experiences, domestic violence, low socio-economic level, unintended pregnancy, and pregnancy at younger ages; these factors vary among different cultures [7,8]. Research studies conducted regionally in Turkey on smaller populations report antenatal depression frequency between 27.9%-33.0%[9-11]. Despite this high frequency of antenatal depression and the importance of prenatal care in Turkey, psychological support during this stressful period does not have any standard care procedure. In this study, we intend to determine the frequency of depressive mood during pregnancy and the associated risk factors in order to contribute to preventive care that can be implemented in this area.

Material and Method

This descriptive study was conducted from February to July 2012 at the antenatal clinic of a major maternity hospital in Ankara, Turkey, a semi-urban region. Of the five Gynecology and Obstetrics polyclinics existing in the study hospital, one was randomly selected. Pregnant women within their third trimester visiting this outpatient polyclinic were included each day in the study. 600 pregnant women were included in the study within a period of six months. Pregnant women were divided into two groups as depressed or non-depressed according to their EPDS results. The study was approved by the Ethics Committee and all pregnant women were informed about the study and an informed written consent was obtained.

After collecting the socio-demographic characteristics and fertility features of patients by face-to-face interview using a questionnaire, depressive symptoms were determined with the Edinburgh Postnatal Depression Scale, the most widely used screening instrument used for antenatal depression in research. Although developed by Cox et al. for determination of postnatal depression, the scale is often used during pregnancy as well [12,13]. The scale contains 10 Likert-type questions in total and provides 4 subscale measurements. Total score is from 0-30. For each item, women are asked to select one of four responses that most closely describe how they have felt over the past 7 days.

The scale was adapted to Turkish by Engindeniz et al. in 1996, its validity and reliability was demonstrated by the same team, sensitivity and selectivity was determined as 84.0% and 88.0%, respectively, and the cut-off score was shown to be ≥ 13 [14]. The EPDS was filled out by each patient one in a private and quiet room.

According to sampling volume formulas, the minimum required sample size was calculated as 545 subjects for this study, assuming a depression frequency of 15.0%, accepting an error margin of ± 0.03 and with a reliability of 95.0%.

Women with depressive symptoms were referred to a psychiatrist for further evaluation and treatment.

Statistical analysis was performed using Statistical Package for Social Sciences version 17 (SPSS Inc. Chicago. IL). Differences between two groups were assessed using Chi-square test, Fisher's Exact test, and Mann-Whitney U-test for categorized variables, Student's t-test for continuous variables, and Multiple Logistic Regression Analysis as multidimensional analysis during the statistical evaluations. $p < 0.05$ was considered as significant.

Results

Average pregnancy week of the subjects was 31.81 ± 2.16 (range, 26-36 weeks). One hundred and fifty-three patients out of 600 were found to be depressive (25.5%) in our study group. Socio-demographic findings related to depressive mood are given in Table 1. Lower and advanced age, low educational level

Table 1. Distribution of socio-demographic characteristics of individuals

Features	Depressed n (%)	Normal n (%)	Total n (%)	P
Age				p=0.009
≤19	25 (34.2)	47 (65.3)	72 (12.0)	
20-24	41 (21.5)	150 (78.5)	191 (31.8)	
25-29	48 (26.8)	131 (73.2)	179 (29.8)	
30-34	17 (17.0)	83 (83.0)	100 (16.7)	
≥35	22 (37.9)	36 (62.1)	58 (9.7)	
Education Level				p=0.002
Elementary school or lower	106 (30.3)	244 (69.7)	350 (58.3)	
High school	41 (21.4)	151 (78.6)	192 (32.0)	
University	6 (10.3)	52 (89.7)	58 (9.7)	
Employment Status				p=0.000
Employed	140 (29.3)	337 (70.7)	477 (79.5)	
Unemployed	13 (10.6)	110 (89.4)	123 (20.5)	
Health Insurance				p=0.000
Yes	137 (24.0)	435 (76.0)	572 (95.3)	
No	16 (57.1)	12 (42.9)	28 (4.7)	
Education level of spouse				p=0.007
Elementary school or lower	90 (30.5)	205 (69.5)	295 (49.2)	
High school	52 (22.8)	176 (77.2)	228 (38.0)	
University	11 (14.3)	66 (85.7)	77 (12.8)	
Employment of spouse				p=0.000
Employed	105 (20.2)	416 (79.8)	521 (86.8)	
Unemployed	48 (60.8)	31 (39.2)	79 (13.2)	
Monthly Income (Turkish Lira)				p=0.000
≤3000	66 (47.1)	74 (52.9)	140 (23.3)	
3001-6000	73 (23.2)	241 (76.8)	314 (52.4)	
≥6001	14 (9.6)	132 (90.4)	146 (24.3)	

of the women/spouse, unemployment, lack of health insurance, and lower monthly income were found to be effective on depression rate ($p < 0.01$).

Correlation between fertility features, chronic conditions, and depressive mood is shown in Table 2. Increased total number of pregnancies, live births, unplanned pregnancy, previous prenatal/postnatal depression diagnosis, depression history in immediate family members, and having a diagnosed chronic disease were found to be correlated with high depression frequency ($p < 0.01$).

Familial factors such as lack of a civil marriage, involuntary marriage, living with spouses' families and the increased number of cohabitants, failure in sharing problems with and receiving support of the spouse/close relatives or friends, and discord/uneasiness within the family were found to be correlated with high depression frequency ($p < 0.01$) (Table 3).

In logistic regression analysis, involuntary marriage, living with spouse's relatives, unplanned pregnancy, existence of a diagnosed chronic disease, not being able to share problems with the spouse and/or with immediate family members/friends were determined as the factors increasing depressive mood risk (Table 4).

Discussion

Frequency of depressive mood within the study group was found as 25.5%. Studies conducted in various countries reported an antenatal depression prevalence of 5.6% in Japan [15], 14.2% in Brazil [8], 17.0% in Sweden [16], and 20.0% [17] in the USA. Various studies performed in Turkey reported an antenatal depression frequency between 27.9%-33.1%, similar to

Table 2. Distribution of the reproductive history and chronic conditions of individuals

	Depressed n (%)	Normal n (%)	Total n (%)	P
Number of pregnancy				$p=0.000$
1	36 (17.7)	167 (82.3)	203 (33.8)	
2-3	78 (25.4)	229 (74.6)	307 (51.2)	
≥ 4	39 (43.3)	51 (56.7)	90 (15.0)	
Live Birth				$p=0.004$
0	50 (20.2)	198 (79.8)	248 (41.3)	
1-2	90 (27.8)	234 (72.2)	324 (54.0)	
≥ 3	13 (46.4)	15 (53.6)	28 (4.7)	
Planned pregnancy				$p=0.000$
Yes	46 (10.4)	396 (89.6)	442 (73.7)	
No	107 (67.7)	51 (32.3)	158 (26.3)	
Depression History				$p=0.000$
Yes	62 (74.7)	21 (25.3)	83 (13.8)	
No	91 (17.6)	426 (82.4)	517 (86.2)	
Previous prenatal/ postnatal depression diagnosis				$p=0.000$
Yes	28 (87.5)	4 (12.5)	32 (8.8)	
No	125 (37.5)	208 (62.5)	333 (91.2)	
Depression history in first degree relatives				$p=0.000$
Yes	71 (68.9)	32 (31.1)	103 (17.2)	
No	82 (16.5)	415 (83.5)	497 (82.8)	
Diagnosed Chronic Disease				$p=0.000$
Yes	56 (44.4)	70 (55.6)	126 (21.0)	
No	97 (20.5)	377 (79.5)	474 (79.0)	

Table 3. Distribution of family features of individuals

Features	Depressed n (%)	Normal n (%)	Total n (%)	P
Legally valid marriage				$p=0.000$
Yes	140 (23.9)	445 (76.1)	585 (97.5)	
No	13 (86.7)	2 (13.3)	15 (2.5)	
Voluntary Marriage				$p=0.000$
Yes	101 (18.7)	440 (81.3)	541 (90.2)	
No	52 (88.1)	7 (11.9)	59 (9.8)	
Number of individuals living at home				$p=0.000$
2	20 (11.4)	155 (88.6)	175 (29.2)	
3-4	74 (24.7)	226 (75.3)	300 (50.0)	
≥ 5	59 (47.2)	66 (52.8)	125 (20.8)	
Association of individuals at home				$p=0.000$
Elementary family	62 (14.6)	362 (85.4)	424 (70.7)	
With spouse's relatives*	84 (59.6)	57 (40.4)	141 (23.5)	
With own relatives*	7 (20.0)	28 (80.0)	35 (5.8)	
Sharing problems with spouse				$p=0.000$
Always	30 (7.6)	363 (92.4)	393 (65.5)	
Sometimes	47 (41.2)	67 (58.8)	114 (19.0)	
Rarely	76 (81.7)	17 (18.3)	93 (15.5)	
Discord/uneasiness within the family				$p=0.000$
Always	48 (100.0)	0 (0.0)	48 (8.0)	
Sometimes	61 (52.6)	55 (47.4)	116 (19.3)	
Rarely	44 (10.1)	392 (89.9)	436 (72.7)	
Sharing problems with relatives/friends				$p=0.000$
Always	23 (7.3)	293 (92.7)	316 (52.7)	
Sometimes	41 (27.5)	108 (72.5)	149 (24.8)	
Rarely	89 (65.9)	46 (34.1)	135 (22.5)	

*In addition to spouse and other children

the results of this study [9-11]. These figures are quite high compared to the rest of the world, which are also affected by factors such as getting pregnant at very young ages and even during adolescence, short intervals between pregnancies, economic challenges, low education levels, family problems, and lack of social support.

Among the study results, the effect of socio-demographic factors on depressive mood is significant which is consistent with previous studies where younger-older age, low education level, unemployment, and lack of health insurance were found to be significant predictors of clinical depression [6,7,16]. It can be said that pregnant women tend to get away from the submissive and desperate approach as their educational level increases and they actively participate in work life and education and employment, all of which help in dealing with stress. Social and economic power, better communication and overall supportiveness of the spouses, and higher educational levels are believed to have a positive impact on depression. Association between psychological disorders and economic income is already known. Many studies report that as the income level decreases, the frequency of antenatal and postnatal depression increases, and income level is a significant risk factor for antenatal depression [6,18]. Lack of partner's support, weak communication and lack of harmony with the partner, and marital problems are found to be correlated with pregnancy and depression and marital problems are indicated to be a potential risk factor for prenatal depression [18].

Table 4. The results for the factors associated with depression in logistic regression analysis

Risk Factors	p	OR	%95 CI
Association of individuals at home			
Elementary family			
With spouse's relatives*	0.706	0.779	0.212-2.860
With own relatives*	0.000	3.731	1.881-7.401
Voluntary marriage			
Yes			
No	0.002	5.642	1.879-16.943
Planned pregnancy			
Yes			
No	0.000	6.348	3.207-12.566
Chronic disease			
No			
Yes	0.001	3.310	1.632-6.712
Sharing problems with spouse			
Always/often			
Sometimes	0.000	3.672	1.784-7.556
Rarely/never	0.000	16.708	6.949-40.175
Sharing problems with relatives / friends			
Always/often			
Sometimes	0.008	2.755	1.297-5.855
Rarely/never	0.000	4.445	2.004-9.860

OR: odds ratio; 95%CI: 95% confidence intervals. *In addition to spouse and other children

Depressive mood rates were found to be higher in women living with their spouses' families and depression rates increase as the number of cohabitants increase. In the literature it is indicated that antenatal depression is less frequently observed among pregnant women living with their spouses in a harmonious marriage and that depression frequency increases when there are more cohabitants who are not part of the elementary family [9]. The group with the highest depressive mood risk is pregnant women who live in the same house with their mothers-in-law and who indicate that their relation is uneasy with a non-supportive mother-in-law [19,20]. In our society, living with the spouse's family is quite common, as this factor has a high percentage within the study group as well. Depression frequency does not increase among pregnant women who live with their own families, which could be related to better communication and sharing levels with their own relatives.

A significant correlation was found between existence of a previous antenatal/postnatal depression diagnosis and observed depression. Moreover, having immediate family members diagnosed with depression also increases depression risk. Depression is known to be a disorder with genetic predisposition, and the existence of depression cases within the family increases depression risk; similar to the results we obtained in this study, there are many research studies in the literature reporting an increase in depression among pregnant women who have family members with a depression history [19,20]. In general, these research studies have concluded that diagnosis of depression or any other psychiatric disorder sometime in the past constitutes a risk factor for antenatal depression, similar to our findings in this study.

An association between chronic diseases and psychiatric disorders, especially depression, has been previously reported [21]. Similarly, studies conducted in Brazil with 326 pregnant subjects with chronic diseases and 2398 subjects with chronic high blood pressure revealed high depression rates, consistent with the results of this study [22,23].

Failure in sharing problems with and receiving support of the spouse, close relatives, or friends significantly increases depression scores. Many studies have shown that depression rates are considerably lower among pregnant women who indicate they have a good relationship and can easily share their problems with their spouses, close relatives, or friends [7,12,18]. It is already known that women can adapt to the concept of maternity more easily in marriages where women can comfortably share significant problems with their spouses and where spouses approve of the maternity role. However women are more inclined to depression in marriages with a poor communication level where the husband fails to show interest and care during and after pregnancy [18].

An association between an unplanned pregnancy and depression during pregnancy has been well-documented [8-10,18]. In a study conducted in Australia on 8556 pregnant women during pregnancy and 6 months after pregnancy, observance of antenatal depression was reported to be higher among subjects with unplanned pregnancy. However, depression was regressed during the postnatal follow-up period [24].

Despite a few studies in the literature showing that the increase in number of pregnancies and births is not a risk factor for antenatal depression and that, in fact, multiparity has a positive protective effect [25], the general view is the existence of a positive relation between number of pregnancies and depression during pregnancy, which is also consistent with the results of this study [7,11]. The increase in number of births and living children also increases the burden on women's shoulders and tires them both physically and psychologically and thus may prepare the grounds for antenatal depression.

Despite this high frequency depression levels observed in Turkey and the importance of the issue, no relevant approach has yet to be developed. Depression must be given a high priority and diligently checked during prenatal healthcare services and diagnosis and treatment approaches should be developed accordingly. All risk factors influencing the development and progress of depression should be carefully interpreted, policies versatile in terms of protection should be introduced, and correct messages should be addressed to society. Among preventive approaches countering depression, priority should be given to increasing women's education levels so women will be provided with employment opportunities where they can develop themselves and ensure their economic independence. Also, income levels, a significant determinant of health, should be improved in line with the foregoing and health insurance and social security should be ensured for everyone. Social awareness should be raised against early and involuntary marriages and common traditional perspectives should be corrected. Men particularly should be educated to improve their approach to and perception of women. Since depression frequency among women increases as the number of pregnancies and births increases, it should be ensured that families have children through conscious choices,

by evaluating the present socio-economic conditions and without neglecting the potential mother's choices and desires. Family planning services should be accessible and approaches that could guide the society in the wrong way should be avoided. Particularly, the negative characteristics of traditional family type which trivializes women should be changed and social awareness should be raised regarding this issue. Depression scanning should be introduced to routine prenatal care programs, it should be possible to make the required interventions, and accessible diagnosis and treatment services should be in place for pregnant women diagnosed with depression. There must be psychiatry units in healthcare centers providing prenatal care programs, due to the fact that the pregnant women, considered to experience depression according to the scale applied in our study, were observed not to apply to relevant healthcare centers for treatment.

The cross-sectional nature of this study and the use of the EPDS screening tool, which is not a diagnostic tool, are limitations of this study in interpreting the results. Large and prospective studies are needed to clarify the risk factors that can cause depression in the antenatal period.

Disclosure of potential conflicts: The authors (Elif Yılmaz, Meltem Çöl, Yasemin Yavuz) have declared no financial or other conflicts of interest.

Competing interests

The authors declare that they have no competing interests

References

- Da Costa D, Larouche J, Dritsa M, Brender W. Psychosocial correlates of prepartum and postpartum depressed mood. *J Affect Disord* 2000;59:31-40.
- Field T, Diego M, Hernandez-Reif M. Prenatal depression effects on the fetus and newborn: a review. *Infant Behav Dev* 2006;29:445-455.
- Dayan J, Creveuil C, Marks MN, Conroy S, Herlicoveiz M, Dreyfus M et al. Prenatal depression, prenatal anxiety, and spontaneous preterm birth: a prospective cohort study among women with early and regular care. *Psychosom Med* 2006;68:938-46.
- Diego MA, Field T, Hernandez-Reif M, Schanberg S, Kuhn C, Gonzalez-Quintero VH. Prenatal Depression restricts fetal growth. *Early Hum Dev* 2009; 85(1), 65-70.
- Beck CT. The effects of postpartum depression on child development: a meta-analysis. *Arch Psychiatr Nurs* 1998;12:12-20.
- Leigh B, Milgrom J. Risk factors for antenatal depression, postnatal depression and parenting stress. *BMC Psychiatry* 2008;16:24.
- Ali NS, Azam IS, Ali BS, Tabbusum G, Moin SS. Frequency and associated factors for anxiety and depression in pregnant women: a hospital-based cross-sectional study. *Scientific World Journal* 2012; 653098:1-9.
- Pereira PK, Lovisi GM, Pilowsky DL, Lima LA, Legay LF. Depression during pregnancy; prevalence and risk factors among women attending a public health clinic in Rio de Janeiro, Brazil. *Cad Saude Publica* 2009;25(12): 2725-2736.
- Çalışkan D, Öncü B, Köse K, Ocaktan ME, Özdemir O. Depression scores and associated factors in pregnant and non-pregnant women: a community-based study in Turkey. *J Psychosom Obstet Gynaecol* 2007; 28:195-200.
- Karaçam Z, Ancel G. Depression, anxiety and influencing factors in pregnancy: a study in a Turkish population. *Midwifery* 2009;25:344-56.
- Şentürk V, Abas M, Berksun O, Stewart R. Social support and antenatal depression in extended and nuclear family environments in Turkey: a cross-sectional survey. *BMC Psychiatry* 2011;11:48.
- Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry* 1987;150:782-2.
- Zhong Q, Gelaye B, Rondon M, Sacnhez SE, Garcia PJ, Sanchez E et al. Comparative performance of Patient Health Questionnaire-9 and Edinburgh Postnatal Depression Scale for screening antepartum depression. *J Affect Disord* 2014;162:1-7.
- Engindeniz A, Küey L, Kültür S. Validity and reliability of Turkish form of Edinburgh Postpartum Depression Scale. *Spring Symposium 1. Book (Turkish)*;1996. P. 51-52.
- Kitamura T, Yoshida K, Okano T, Kinoshita K, Hayashi M, Toyoda N et al. Multicentre prospective study of perinatal depression in Japan: incidence and

correlates of antenatal and postnatal depression. *Arch Womens Ment Health* 2006;9(3):121-30.

- Josefsson A, Berg G, Nordin C, Sydjo G. Prevalence of depressive symptoms in late pregnancy and postpartum. *Acta Obstet Gynecol Scand* 2001;80:251-255.
- Marcus SM, Flynn HA, Blow FC, Barry KL. Depressive symptoms among pregnant women screened in obstetrics settings. *J Womens Health* 2003;12(4):373-80.
- Lancaster CA, Gold KJ, Flynn HA, Yoo H, Marcus SM, Davis MM. Risk factors for depressive symptoms during pregnancy: a systematic review. *Am J Obstet Gynecol* 2010;202:5-14.
- Lau Y, Keung DW. Correlates of depressive symptomatology during the second trimester of pregnancy among Hong Kong Chinese. *SocSci Med* 2007;64:1802-1811.
- Johnstone SJ, Boyce PM, Hickey AR, Morris-Yatees AD, Harris MG. Obstetric risk factors for postnatal depression in urban and rural community samples. *Aust N Z J Psychiatry* 2001;35:69-74.
- Wells KB, Golding JM, Burnam MA. Psychiatric disorder in a sample of the general population with and without chronic medical conditions. *Am J Psychiatry* 1988;145(8):976-81.
- Benute GRG, Nomura RMY, Reis JS, Junior RF, Souza De Lucia MC, Zugaib M. Depression during pregnancy in women with a medical disorder: risk factors and perinatal outcomes. *Clinics* 2010;65:1127-1131.
- Katon WJ, Russo JE, Melville JL, Katon JG, Gavin AR. Depression in pregnancy is associated with preexisting but not pregnancy-induced hypertension. *Gen Hosp Psychiatry* 2012;34(1):9-16.
- Najman JM, Morrison J, Williams G, Andersen M, Keeping JD. The mental health of women 6 months after they give birth to an unwanted baby: alongitudinal study. *Soc Sci Med* 1991;32:241-7.
- Barbadoro P, Cotichelli G, Chiatti C, Simonetti ML, Marigliano A, Di Stanislao F et al. Socio-economic determinants and self-reported depressive symptoms during postpartum period. *Women Health* 2012;52:352-68.

How to cite this article:

Yılmaz E, Col M, Yavuz Y. The Frequency of Depressive Mood and Associated Factors in Pregnant Women in a Semi-Urban Region. *J Clin Anal Med* 2017;8(suppl 4): 261-5.