



BODY MASS INDEX RELATED CLINICAL OUTCOMES AFTER KNEE REPLACEMENT IN OVERWEIGHT AND OBESE PATIENTS

AŞIRI KİLOLU VE OBEZ HASTALARDA DİZ PROTEZİ SONRASI KLİNİK SONUÇLARIN VÜCUT KİTLE İNDEKSİ İLE İLİŞKİSİ

BODY MASS INDEX RELATED CLINICAL OUTCOMES AFTER KNEE REPLACEMENT

Emre Bilgin¹, Hasan Bombacı², Ali Turgut¹, Önder Kalenderer¹, Bekir Eray Kılıç³, Levent Adıyeko²

¹Department of Orthopaedics and Traumatology, Health Science University Tepecik Training and Research Hospital, İzmir,

²Department of Orthopaedics and Traumatology, Health Science University Haydarpaşa Numune Training and Research Hospital, İstanbul,

³Department of Orthopaedics and Traumatology, Gölhisar State Hospital, Burdur, Turkey

Öz

Amaç: Aşırı kilolu ve obez hastalarda total diz protezi (TDP) sonrası vücut kitle indeksinin (VKİ) fonksiyonel sonuçlara etkisini ortaya koymak. Gereç ve Yöntem: Ocak 2002 ve Aralık 2010 arasında primer TDP yapılan hastalar geriye dönük incelendi. Ortalama takip süresi 7 yıldır. Dizler vücut kitle indeksleri baz alınarak Dünya Sağlık Örgütü sınıflamasına göre gruplara ayrıldı ve aşırı kilolu, 1. derece obez ve 2. derece obez hastalar değerlendirildi. Grupların klinik sonuçları Oxford Diz Skoru (OKS), Görsel ağrı skalası (VAS) ve Kısa Form-36 fiziksel skoru (SF-36 PCS) ile Kısa Form-36 mental skoru (SF-36 MCS) hesaplanarak karşılaştırıldı. Bulgular: En düşük skorlar 2. derece obez hastalarda elde edildi. Fakat gruplar arasında OKS, SF-36 PCS, SF-36 MCS ve VAS skorları arasında istatistiksel olarak anlamlı farka rastlanmadı. Grupların ortalama yaşları sırasıyla 67, 64 ve 62 idi ($p < 0.05$). Diğerleri ile karşılaştırıldığında 2. derece obez grubundaki hastaların daha erken yaşta ameliyat edildiği görüldü. Tartışma: Aşırı kilolu, 1. derece obez ve 2. derece obez hastalarda TDP sonrası klinik sonuçların ortalama 7 yıllık takipte VKİ'nden etkilendiği görüldü.

Anahtar Kelimeler

Artroplastisi; Diz; Vücut Kitle İndeksi; Klinik Sonuç; Fonksiyonel Skor

Abstract

Aim: To analyze the influence of body mass index (BMI) on functional outcome in overweight and obese patients after total knee replacement (TKR). Material and Method: The patients who underwent primary TKR between January 2002 and December 2010 were retrospectively reviewed. The mean follow-up period was seven years. The knees were divided into the groups according to the World Health Organization classification based on BMIs and overweight, class 1 obese, class 2 obese patients were assessed. The clinical outcomes of the groups were compared by evaluating the Oxford Knee Score (OKS), visual analog scale (VAS), and Short Form-36 physical component summary (SF-36 PCS) and mental component summary (SF-36 MCS) scores. Results: The lowest scores were obtained in class 2 obese group. However, the differences of OKS, SF-36 PCS, and SF-36 MCS and VAS between the groups were not found to be statistically significant. The mean ages of the groups were 67, 64 and 62 respectively ($p < 0.05$). Class 2 obese group underwent surgery at a younger age compared to the others. Discussion: Clinical outcome after TKR was not affected by BMI at a mean seven years follow-up in overweight, class 1 obese and class 2 obese patients.

Keywords

Arthroplasty; Knee; Body Mass Index; Clinical Outcome; Function Score

DOI: 10.4328/JCAM.5123

Received: 03.06.2017 Accepted: 14.06.2017 Published Online: 15.06.2017

Corresponding Author: Emre Bilgin, Department of Orthopaedics and Traumatology, Health Science University Tepecik Training and Research Hospital, İzmir, Turkey. T.: +90 2324696969-1415 GSM: +905067878430 F.: +90 2324330756 E-Mail: dremrebilgin@hotmail.com

Introduction

Obesity is a significant health condition associated with multiple comorbidities, including diabetes mellitus, metabolic syndrome, hypertension and coroner artery diseases [1]. The prevalence of obesity has been increasing in both emerging and industrialized countries [2]. It was stated that a 33% increase in obesity prevalence and a 130% increase in severe obesity prevalence are estimated over the next two decades [3].

Obesity is also defined as a risk factor for the development of hip and knee osteoarthritis [4-6]. Recently, the prevalence of overweight and obese individuals has been increasing among orthopedic patients who underwent total knee replacement (TKR) [7]. Furthermore, need for having TKR has been increasing due to increased body mass index (BMI) [8]. Thus, it is not surprising that obese patients are undergoing surgery at a younger age compared with their non-obese counterparts [9,10].

The current literature has divided in opinion of the influence of BMI on clinical outcomes after TKR. Several studies reported that obesity adversely affected the functional scores [11-14]. Others claimed that the effect of BMI on clinical outcomes was not significant after TKR [15-17]. Most of them compared the obese (BMI>30) and non-obese (BMI<30) individuals [13,16,17], or intercalarily morbidly obese (BMI>40) patients [11,12,17]. However, only a few studies have reported subgroup analysis by classifying the overweight and obese patients as recommended by World Health Organization (WHO) [10,18].

The aim of this study was to determine the influence of body mass index on functional outcome among overweight and obese patients after TKR. It was hypothesized that higher BMI is associated with poorer functional scores.

Material and Method

The study was approved by the local Ethics Committee. The patients who underwent primary TKA in a single clinic between January 2002 and December 2010 were reviewed retrospectively. The inclusion criteria for the study were patients with primary osteoarthritis who underwent primary TKA at least five years before their most recent follow-up visit and overweight or obese patients according to their BMIs. Patients who were over 80 years old on their last visit, had prior knee surgery or had <5-year follow-up period were excluded.

The clinicodemographic data including sex, age, operation side and follow-up periods were collated for each patient. The BMIs were calculated for each patient as body weight(kg)/height²(m²) at their most recent follow-up visit, then Function scores and life-quality assessment were obtained. Function assessment was performed using the Oxford Knee Score (OKS). To determine the relief of pain and quality of life, the visual analog scale (VAS) and Short Form 36 (SF-36) health survey questionnaire—both physical component summary (PCS) and mental component summary (MCS)—were used. In bilateral cases, each knee was included into the belonged group according to the BMI. OKS and VAS which reflect the related knee were measured separately for each knee. However, SF-36 scores which reflect the general condition of the patient were the same for both of the knees.

The knees were divided into the groups according to the WHO classification based on BMIs as overweight (25 to 29.99), class

1 obese (30 to 34.99), class 2 obese (35 to 39.99) and class 3 obese (>40) [19]. The function scores of each group were compared using the OKS, VAS, SF-36 PCS, and SF-36 MCS.

Surgical technique

All operations were performed using a conventional approach with applying a tourniquet by three different senior orthopedic surgeons. A standard surgical approach with a medial parapatellar incision and arthrotomy was used. Femoral cuts were performed by intramedullary guides, and extramedullary guides were used for tibial cuts in all cases. Cemented cruciate ligament retaining modern implant designs were used without patellar resurfacing. All patients received antibiotics prophylaxis for 24 hours and, low molecular weight heparin and full-length anti-embolic stockings were applied for deep venous thromboembolism prophylaxis after surgery. The patients were mobilized the next day after surgery, and all patients underwent standardized rehabilitation programs.

Statistical Analysis

Statistical analysis was performed using IBM SPSS Statistics for Windows, version 24.0. Armonk, NY: IBM corp., USA. Different categorical data were compared using the chi-square test and Fisher–Freeman–Halton test. The results of normal distributions were compared using one-way ANOVA test, and non-normal distributions were compared using the Kruskal–Wallis test. Statistical significance was set as $p < 0.05$.

Results


Sufficient data of 237 patients were obtained (preoperative clinicodemographic data, phone number, etc.) from the hospital archives who had proper follow up. Fifty-three of them were over 80 years old at the last follow-up, and 17 of them died. Eighteen patients underwent surgery for seconder osteoarthritis etiology, and 16 of the patients had prior knee surgery. These patients were excluded according to the inclusion and exclusion criteria. Thirty-five patients stated that they had no complaint and did not want to continue follow-ups. The other 45 patients rejected to participate in the study. Remaining 53 patients were included in the survey.

Because there were only two morbidly obese (class 3 obese) patients and the number was not sufficient for statistical comparison, these patients were excluded. A Total of 70 knees of 51 patients were evaluated, and the comparison was performed between overweight (n=22 knees), class 1 obese (n=25 knees) and class 2 obese (n=23 knees) patients. The distribution of the groups based on BMIs are shown in Table 1.

There were no complications, such as wound problem, deep infection, deep venous thrombosis, periprosthetic fracture, or component loosening.

The mean age of the patients at surgery date was 64.8 (range; 56–73), and mean follow-up period was 84.9 months (range; 60–144). The mean ages of the groups were 67, 64 and 62 respectively. The difference in the ages between the groups was statistically significant. It was demonstrated that obese patients underwent surgery at a younger age compared to overweight patients as estimated. On the other hand, the differences were not significant between the groups according to the

Table 1. Distribution of the number of the groups based on BMIs



follow-up periods, gender distribution, or operation sides. The clinicodemographic data of each group was shown in Table 2. The mean OKS of the groups were 35.8, 37.1 and 33.5 respectively ($p>0.05$). The mean VAS of each group was 2.9, 2.8 and 3.8 respectively ($p>0.05$). The mean SF-36 PCS of the overweight group was 38.5, and class 1 obese group was 39.9. Class 2 obese group had the lowest SF-36 PCS (35.9). However, the differences were not statistically significant. The SF-36 MCSs of the groups were 46.2, 45.4 and 44.4 respectively ($p>0.05$). The differences of OKS, VAS, SF-36 PCS and SF-36 MCS between the groups were not found to be statistically significant. The function scores of each group were shown in Table 3.

Discussion

The overall results of the current study support that clinical outcome after TKR is not related to the BMI in overweight and obese individuals at mid-term.

Table 2. Clinicodemographic data of groups

	Overweight	Class 1 Obese	Class 2 Obese	P value
Mean age	67±4.1	64.7±3.3	62.8±3.7	^a 0.02
Mean BMI	27.3±1.3	32.2±1.4	37.9±1	^b <0.001
Mean Follow-up period (month)	90±13	80.9±13.6	84.5±19.8	^b 0.68
	n	n	n	
Gender	Female	19	21	^c 0.14
	Male	3	4	
Operation side	Right	11	13	^d 0.90
	Left	11	12	

^aOne way ANOVA Test, ^bKruskal Wallis Test, ^cFisher Freeman-Halton Test, ^dPearson Chi-Square Test

Table 3. Function scores of the groups

	Overweight	Class 1 Obese	Class 2 Obese	P value
OKS (Mean±Sd)	35.8±7.7	37.1±8.2	33.5±7.4	^b 0.93
VAS (Mean±Sd)	2.9±2.3	2.8±1.8	3.8±2.1	^b 0.14
SF-36 PCS (Mean±Sd)	38.5±10	39.9±9.7	35.9±9.1	^b 0.33
SF-36 MCS (Mean±Sd)	46.2±10.1	45.4±11.5	44.4±11	^a 0.85

^aOne-way ANOVA Test, ^bKruskal Wallis Test

There is a debate in the literature regarding the influence of BMI on clinical outcomes after TKR. Several short-term studies showed no difference in clinical outcomes between obese and non-obese individuals [20,21]. Other studies with a mid-term or longer follow-up were also unable to demonstrate inferior results in obese patients [15,16,22].

However, Foran et al. reported that obese patients with BMI >30 showed inferior Knee Society Score and higher revision rates at 15 years after TKR [13]. Similarly, Jackson et al. found significantly lower post-operative total clinical scores in obese patients after cementless TKR at mean 9.2 years [14]. On the other hand, a greater decrease in clinical scores was reported in the morbidly obese group, and these patients were advised to lose weight before the surgery [11].

To our knowledge only one study compared the clinical outcomes after TKR by classifying the patients as overweight, class 1 obese, class 2 obese and class 3 obese like our study method as recommended by WHO. O'Neill et al. reviewed 2180 TKR, and in contrast to our results, they demonstrated significantly inferior scores in class 2 obese patients compared to both overweight and class 1 obese patients [18]. Although a small number of the patients limits the power of our study, the results of our study are more reliable when considering our mid-term (84 months=7 years) follow-up period compared to their only mean 19 months follow-up.

Collins et al. also evaluated 445 TKR with considering the overweight patients in non-obese group and dividing the obese group as mildly obese (BMI 30 to 35) and highly obese (BMI>35), then they reported significantly poorer results in highly obese group compared to non-obese and mildly obese group [23]. They used only KSS for clinical evaluation, whereas multiple scores including OKS, VAS, and SF-36 were used for evaluation of clinical outcomes in our study.

The patients who had had a higher BMI needed TKR at a younger age in our study. Similarly, Guanter et al. reported that the overweight, class 1 obese and class 2 obese patients underwent total joint replacement surgery at the mean age of 68, 65 and 64 respectively [10]. Thus, can also provide the evidence that obesity is a risk factor for the development of knee osteoarthritis [4-6].

The major limitation of the current study is its retrospective design. The second limitation can be stated as our mid-term follow-up period. This is particularly important about the development of aseptic loosening, implant failures and subsequent need for revision surgery. Although similar revision rates and good clinical outcomes were reported in both obese and non-obese patients in 10 years follow-up, there is doubt in longer period results [22].

This study only included the obese patients with BMIs between 30 and 40. Our results would be more reliable and powerful whether we could perform the statistical comparison of class 3 obese patients with BMIs over 40. Nevertheless, we revealed that TKR can be carried out in obese whose BMIs between 30 and 40 with results similar to those performed in overweight individuals.

In summary, it was demonstrated that clinical outcome after TKR is not related to patient's BMI at mean seven years follow-up. We recommend starting diet programs with the surgery, instead of denying the obese patients for TKR.

Funding

The funders had no role in study design, data collection, and analysis, decision to publish, or preparation of the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest regarding the submission and publication of this manuscript

Animal and human rights statements

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

References

- Allen SR. Total knee and hip arthroplasty across BMI categories: a feasible option for the morbidly obese patient. *J Surg Res* 2012; 175(2): 215-7
- Campos P, Saguy A, Ernsberger P, Oliver E, Gaesser G. The epidemiology of overweight and obesity: public health crisis or moral panic? *Int J Epidemiol* 2006; 35(1): 55-60.
- Finkelstein EA, Khavjou OA, Thompson H, Trogdon JG, Pan L, Sherry B et al. Obesity and severe obesity forecasts through 2030. *Am J Prev Med* 2012; 42(6): 563-70
- Stürmer T, Günther KP, Brenner H. Obesity, overweight and patterns of osteoarthritis: The Ulm Osteoarthritis Study. *J Clin Epidemiol* 2000; 53(3): 307-13.
- Richmond SA, Fukuchi RK, Ezzat A, Schneider K, Schneider G, Emery CA. Are joint injury, sport activity, physical activity, obesity, or occupational activities predictors for osteoarthritis? A systematic review. *J Orthop Sports Phys Ther* 2013; 43(8): 515-B19
- Hootman JM, Helmick CG. Projections of US prevalence of arthritis and associated activity limitations. *Arthritis Rheum* 2006; 54(1): 226-9
- Woon CY, Piponov, H, Schwartz BE, Moretti, VM, Schraut NB, Shah RR et al. Total Knee Arthroplasty in Obesity: In-Hospital Outcomes and National Trends. *J Arthroplasty* 2016; 31(11): 2408-14.
- Bourne R, Mukhi S, Zhu N, Keresteci M, Marin M. Role of obesity on the risk for total hip or knee arthroplasty. *Clin Orthop Relat Res* 2007; 465: 185-8
- Changulani M, Kalairajah Y, Peel T, Field RE. The relationship between obesity and the age at which hip and knee replacement is undertaken. *J Bone Joint Surg Br* 2008; 90(3): 360-3.
- Guenther D, Schmidl S, Klatte TO, Widhalm HK, Omar M, Krettek C et al. Overweight and obesity in hip and knee arthroplasty: Evaluation of 6078 cases. *World J Orthop* 2015; 6(1): 137-44
- Amin AK, Clayton RA, Patton JT, Gaston M, Cook RE, Brenkel IJ. Total knee replacement in morbidly obese patients: results of a prospective, matched study. *J Bone Joint Surg Br* 2006; 88(10): 1321-6.
- Dowsey MM, Liew D, Stoney JD, Choong PF. The impact of pre-operative obesity on weight change and outcome in total knee replacement: a prospective study of 529 consecutive patients. *J Bone Joint Surg Br* 2010; 92(4): 513-20.
- Foran JR, Mont MA, Etienne G, Jones LC, Hungerford DS. The outcome of total knee arthroplasty in obese patients. *J Bone Joint Surg Am* 2004; 86-A (8): 1609-15.
- Jackson MP, Sexton SA, Walter WL, Walter WK, Zicat BA. The impact of obesity on the mid-term outcome of cementless total knee replacement. *J Bone Joint Surg Br* 2009; 91(8): 1044-8.
- Amin AK, Patton JT, Cook RE, Brenkel IJ. Does obesity influence the clinical outcome at five years following total knee replacement for osteoarthritis? *J Bone Joint Surg Br* 2006; 88(3): 335-40.
- Hamoui N, Kantor S, Vince K, Crookes PF. Long-term outcome of total knee replacement: does obesity matter? *Obes Surg* 2006; 16(1): 35-8.
- Ersozlu S, Akkaya T, Ozgur AF, Sahin O, Senturk I, Tandoğan R. Bilateral staged total knee arthroplasty in obese patients. *Arch Orthop Trauma Surg* 2008; 128(2): 143-8.
- O'Neill SC, Butler JS, Daly A, Lui DF, Kenny P. Effect of body mass index on functional outcome in primary total knee arthroplasty - a single institution analysis of 2180 primary total knee replacements. *World J Orthop* 2016; 7(10): 664-9.
- BMI classification. http://apps.who.int/bmi/index.jsp?introPage=intro_3.html; 2017 25/02/2017.
- Stickless B, Phillips L, Brox WT, Owens B, Lanzer W. Defining the relationship between obesity and total joint arthroplasty. *Obes Res* 2001; 9(3): 219-23.
- Deshmukh RG, Hayes JH, Pinder IM. Does body weight influence outcome after total knee arthroplasty?: A 1-year analysis. *J Arthroplasty* 2002; 17(3): 315-9.
- Griffin FM, Scuderi G, Insall J, Colizza W. Total knee arthroplasty in patients who were obese with 10 years follow-up. *Clin Orthop* 1998; 356:28-33.
- Collins RA, Walmsley PJ, Amin AK, Brenkel IJ, Clayton RAE. Does obesity influence clinical outcome at nine years following total knee replacement? *J Bone Joint Surg Br* 2012; 94(10): 1351-5.

How to cite this article:

Bilgin E, Bombacı H, Turgut A, Kalenderer Ö, Kılınc BE, Adıyke L. Body Mass Index Related Clinical Outcomes After Knee Replacement in Overweight and Obese Patients. *J Clin Anal Med* 2017; DOI: 10.4328/JCAM.5123.